

# Johne's Disease Vaccines Past, Present and Future



**Shanmugasundaram K and Bhupendra Nath Tripathi\***

National Centre for Veterinary Type Cultures-National Research Centre on Equines, India

Submission: June 23, 2018; Published: August 24, 2018

\*Corresponding author: Bhupendra Nath Tripathi, National Centre for Veterinary Type Cultures-National Research Centre on Equines, Sirsa Road, Hisar-125001, Haryana, India, Tel: +91-1662275787; Email: [bntripathi1@yahoo.co.in](mailto:bntripathi1@yahoo.co.in)

## Abstract

Control and eradication of Johne's disease of ruminants caused by *Mycobacterium aviumsubsp paratuberculosis* (MAP) still remains a formidable challenge before the scientific community because of insidious nature of the infection and our inability to diagnose subclinical cases. Vaccination against MAP is considered an alternate option to control JD. Inactivated vaccines, live attenuated vaccines, subunit vaccines, DNA vaccines and recombinant protein based vaccines have been developed for JD with variable successes. In this mini review, we have discussed various types of vaccines developed of JD and their current status. Searching of new vaccines and testing against MAP infection is still a very practical approach to control and eradicate the JD.

**Keywords:** Disease Vaccines; MAP; Johne's Disease; live attenuated vaccines; *Mycobacterium aviumsubsp paratuberculosis*

**Abbreviations:** LAV: Live Attenuated Vaccines; MAP: *Mycobacterium aviumsubsp paratuberculosis*; JD: Johne's Disease

## Introduction

Johne's Disease (JD) or paratuberculosis is an incurable, chronic granulomatous enteritis of ruminants and other animals and is caused by *Mycobacterium aviumsubsp paratuberculosis* (MAP) [1]. The organism has also been implicated with Crohn's disease of humans. JD incurs huge economic losses to the dairy and small ruminant industry worldwide. During early stages of the infection, infected animals shed MAP intermittently in the feces and thus acts as a source of infection to the susceptible animals [2]. Lack of proper diagnostic assays in identification of the subclinical cases of MAP infection is a major challenge in the control and eradication of the disease [3]. Alternatively, vaccination against MAP is a viable option to control and reduce the economic burden due to JD [3]. In this mini review, we have discussed the recent development about vaccination against MAP infection and futurology.

Vaccination trial against MAP infection was initiated in 1990 with whole cell antigens obtained from heat killed MAP organisms and was suspended in oil adjuvants [4]. Inactivated whole MAP cells have been used to prepare commercial vaccines such as Mycopar, Gudair and Silirum. Mycopar was developed from MAP strain 18 (now identified as *M. avium* subsp *avium*), whereas Gudair and Silirum from MAP F316 strain [5]. These vaccines have been shown to minimize MAP shedding and disease transmission but do not protect the animals from new infections [6].

Sequence analysis and annotation of MAP genomes, prediction of proteins (structure, sub-cellular locations and antigenicity) based on the bioinformatics analysis and subsequent validation with laboratory experiments opened up a new era in the development of vaccines against MAP [6,7]. A rational frame model has been proposed recently to test the new generation vaccine against MAP. It consists of three phases such as phase I (screening of candidates in bovine macrophages), phase II (mouse challenge models) and phase III (goat challenge model) to develop MAP vaccines [8].

## Vaccine Types

Presently live attenuated vaccines (LAV), subunit vaccines, DNA vaccines and recombinant protein based vaccines are available for the JD with variable success reports. Several mutants of MAP have been produced by allelic exchange, phage-mediated and transposon mutagenesis to attenuate the MAP virulence [3]. Recently, transposon mutant bank was constructed to 13,536 MAP K-10 Tn5367 and some of the mutants like 4H2, 30H9, 22F4 and 40A9 showed reduced virulence with Bovine macrophages and Monocyte-derived macrophages cells [9]. However, further studies are warranted to assess their suitability as LAV candidates. Mutants generated by direct allelic exchange methods targeting known virulence genes such as *pknG*, *relA*, *ppiA*, *mpt64*, *lsr2*, *leuD*, *sigL*, *sigH* and *secA2* [3,10,11] were screened as LAV and the results showed that  $\Delta relA$ ,  $\Delta ppiA$ ,  $\Delta leuD$ ,  $\Delta mpt64$ ,  $\Delta secA2$   $\Delta sigL$  and  $\Delta sigH$  had protective immune

response against MAP infection in the experimental models tested. LAV stimulates both innate and adaptive immune response that is considered an advantage with these vaccines. However, it would not differentiate between infected and the vaccinated animals[6].

Subunit vaccines using MAP DNA or recombinant protein antigens have also been tested. These vaccines may overcome the interference in the diagnosis of bovine tuberculosis in comparison to the whole MAP cell based vaccines (killed or attenuated)[3]. Different antigens like antigen 85 Complex (A, B and C), LprG, MAP1518, MAP0261c, MAP2698c, MAP3567, MAP3184, SodD, AhpC, AhpD and Hsp70 were tested as subunit vaccines. These studies were mainly focused on the protective immune responses due to cell mediated immune responses rather than humoral immune responses[3]. However, it has been shown that a protective response from Hsp70 was due to activation of B lymphocytes. Unfortunately, none of subunit vaccine candidates tested so far could be able to provide complete protection in the murine, calf and goat models[8].

Delivery of expressed MAP antigens through attenuated strains like *Salmonella* and *Lactobacillus salivarius* was shown to be an alternate way to stimulate protective mucosal immune responses[12,13]. This approach is, however, in primitive stage and needs further studies to prove its usefulness in the development of subunit vaccines. Combination of viral vectors with MAP antigens was also tried to develop DNA subunit vaccines. *AhpC*, *gsd*, *p12* and *mpa* gene fusion constructs were developed with viral vectors and used to test their immune protective role against MAP. Results showed protection against subsequent challenges studies in murine models[14]. In another study in which MAP antigens were delivered through non-replicative human adenovirus 5 and modified vaccinia virus Ankara recombinant, induced MAP specific CD4<sup>+</sup> and CD8<sup>+</sup> immune responses and protected from the MAP infection [15].

### Conclusions and Future Directions

Diverse strategies have been used to develop vaccines against MAP. Unfortunately, as of now none of the live attenuated or subunit vaccines are available commercially. The most important areas to focus are

- i) MAP cell biogenesis to understand the biology of MAP,
- ii) Host-pathogen interactions to know about how bacteria overcomes highly orchestrated host defence mechanisms such as innate and adaptive immunity,
- iii) Pathogenomic analysis of MAP strains to know geographical distribution and SNPs to select an appropriate MAP strain for further studies,
- iv) Top down proteomic approaches to identify more MAP-specific antigens and their iso-forms for further screening,
- v) Studies on post translational modification of MAP proteins to understand the pathobiology and immunogenicity

and thus will offer to select new level MAP-specific epitopes as a better vaccine candidates and

vi) Long term trials that mimic with natural MAP infection. Even though, as of now none of the vaccines gives complete protection against MAP, but searching of new vaccines and testing against MAP infection is still a very practical approach to control and eradicate the JD.

### References

1. Narnaware SD, Sivakumar P, Tripathi BN (2016) Pathology of classical and non-classical immunopathologic forms of Johne's disease in bullocks naturally infected with *Mycobacterium avium* subspecies *paratuberculosis* bison type. *Research in Veterinary Science* 106: 74-80.
2. Garcia AB, Shalloo L (2015) *Invited review: The economic impact and control of paratuberculosis in cattle.* *J Dairy Sci* 98(8): 5019-5039.
3. Park HT, Yoo HS (2016) Development of vaccines to *Mycobacterium avium* subsp. *Paratuberculosis* infection. *Clin Exp Vaccine Res* 5(2): 108-116.
4. Kormendy B (1994) The effect of vaccination on the prevalence of paratuberculosis in large dairy herds. *Vet Microbiol* 41(1-2): 117-125.
5. Bastida F, Juste RA (2011) Paratuberculosis control: a review with a focus on vaccination. *J Immune Based Ther Vaccines* 9(8): 1-17.
6. Bannantine JB, Talaat AM (2015) Controlling Johne's disease: vaccination is the way forward. *Front Cell Infect Microbiol* 5(2): 1-2.
7. Gurung RB, Purdie AC, Whittington RJ, Begg DJ (2014) Cellular and humoral immune responses in sheep vaccinated with candidate antigens MAP2698c and MAP3567 from *Mycobacterium avium* subspecies *paratuberculosis*. *Front Cell Infect Microbiol* 4: 1-8.
8. Bannantine JP, Hines ME, Bermudez LE, Talaat AM, Sreevatsan S, et al. (2014) A rational framework for evaluating the next generation of vaccines against *Mycobacterium avium* subspecies *paratuberculosis*. *Front Cell Infect Microbiol* 4(126): 1-10.
9. Rathnaiah G, Lamont EA, Harris NB, Fenton RJ, Zinniel DK et al. (2014) Generation and screening of a comprehensive *Mycobacterium avium* subsp. *paratuberculosis* transposon mutant bank. *Front Cell Infect Microbiol* 4(144): 1-17.
10. Ghosh P, Steinberg H, Talaat AM (2014) Virulence and immunity orchestrated by the global gene regulator *sigL* in *Mycobacterium avium* subsp. *paratuberculosis*. *Infect Immun* 82(7): 3066-3075.
11. Ghosh P, Shippy DC, Talaat AM (2015) Superior protection elicited by live-attenuated vaccines in the murine model of paratuberculosis. *Vaccine* 33(51):7262-7270.
12. Faisal SM, Yan F, Chen TT, Useh NM, Guo S, et al. (2013) Evaluation of a *Salmonella* vectored vaccine expressing *Mycobacterium avium* subsp. *paratuberculosis* antigens against challenge in a goat model. *PLoS ONE* 8(8): e70171.
13. Johnston CD, Bannantine JP, Govender R, Endersen L, Pletzer D, et al. (2014) Enhanced expression of codon optimized *Mycobacterium avium* subsp. *paratuberculosis* antigens in *Lactobacillus salivarius*. *Front Cell Infect Microbiol* 4(120):1-10.
14. Bull TJ, Gilbert SC, Sridhar S, et al. (2007) A novel multi-antigen virally vectored vaccine against *Mycobacterium avium* subspecies *paratuberculosis*. *PLoS One* 2(11): e1229.
15. Bull TJ, Vrettou C, Linedale R, McGuinness C, Strain S, et al. (2014) Immunity, safety and protection of an Adenovirus 5 prime: modified Vaccinia virus Ankara boost subunit vaccine against *Mycobacterium avium* subspecies *paratuberculosis* infection in calves. *Vet Res* 45:112.



This work is licensed under Creative Commons Attribution 4.0 License  
DOI: [10.19080/AIBM.2018.10.555784](https://doi.org/10.19080/AIBM.2018.10.555784)

### Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats  
**( Pdf, E-pub, Full Text, Audio )**
- Unceasing customer service

Track the below URL for one-step submission  
<https://juniperpublishers.com/online-submission.php>