

Research Article

Volume 11 Issue 5 - November 2018
DOI: 10.19080/AIBM.2018.11.555823

Adv Biotech & Micro

Copyright © All rights are reserved by Mustafa Riadh Hussien

Toxoplasmosis Seropositivity and Male Sex Hormones



Mustafa Riadh Hussien^{1*}, Adel T M AlSaeed² and Souzan H Eassa³

¹Department of Microbiology, College of Nursing, Iraq

²Department of Microbiology, College of Medicine, Iraq

³Department of Immunoparasitology, HOHED, College of Medicine, Iraq

Submission: April 09, 2018; **Published:** November 16, 2018

***Corresponding author:** Mustafa RiadhHussien, Department of Microbiology, College of Nursing, Iraq

Abstract

Background: Toxoplasmosis is a cosmopolitan disease with acute and chronic infections, caused by the obligate intracellular protozoan parasite *Toxoplasma gondii* that can infect a variety of cells in almost all warm-blooded animals including humans. The study aimed to determine the seroprevalence of *T. gondii* infection among males in Duhok city using ELISA (IgG and IgM).The relationship between toxoplasmosis and reproductive hormones including testosterone, free testosterone, and Follicle Stimulating Hormone (FSH) levels and its association with male sterility were also investigated.

Keywords: Toxoplasmosis; Acute and chronic infection; Male sex hormones

Introduction




Figure 1: Blood groups distribution among males enrolled in the study.

Regarding Iraq and Kurdistan region, there are limited studies in this direction such as, Mahmood SH [1]who studied (the effect of toxoplasmosis on the level of some male sex hormones in samples from National BloodTransfusion Center/Baghdad) and also BassadA. AL-Aboody[2]in Iraq/Thiqar who studied the (the prevalence of toxoplasmosis among male blood donors). Regarding to Kurdistan, In Kalar city, a study by Al-Bajalan et al. [3]indicated that there is a relationship presence of an association between toxoplasmosis and secondary infertility. Since the prevalence of *T. gondii* in primary infertile, secondary infertile and fertile males among 260 persons attending the hospital were 45.16%, 53.33%, and 47.37%, respectively by using ELISA tests,

respectively(Figure 1).In Duhok province, there is no any study dealing with the effect of toxoplasmosis on the levels of male sex hormones and fertility. Therefore, the main objective of the present study is to find out the correlation between *T. gondii* infection (acute, chronic), and the levels of serum testosterone, serum follicle stimulating hormone and fertility in a group of healthy blood donor males and those who visited the IVF center of Azadi teaching hospital[4-20](Table1-2).

Table 1: The distribution of the studied males according to their characteristics (no= 248).

Characteristics of Subjects	Groups	N	%
Residence	Urban Rural	125123	50.4 49.5
Occupation	Official Private	134105	57.6 42.3
Fertility	Fertile Infertile	22226	89.5 10.5
Blood groups	O+	80	32.2
	O-	7	2.8
	A+	57	22.9
	A-	5	2
	B+	68	27.4
	B-	4	1.6
	AB+	25	10
	AB-	2	0.8

Age groups	21-30	63	25.4
	31-40	105	42.3
	41-50	67	27
	51-60	13	5.2

Table 2: Statistical distribution of the hormones (TTH, FTH and FSH) characteristics of males enrolled in the study using ELISA IgG and IgM Abs.

Males' Characteristics	Mean	S. D
Total Testosterone (ng/ml)	6.16	10.3
Free Testosterone (pg.ml)	14.81	4.71
Follicle-stimulating hormone (IU/ml)	4.84	2.2
Anti-toxoplasma IgM	0.541	0.22
Anti-toxoplasma IgG	0.625	0.75

In order to determine

- a. The rate of Toxoplasma infections among males in Duhok city using ELISA (IgM and IgG).
- b. To find out the relationship between toxoplasmosis and blood groups and some demographic factors.
- c. To investigate the relationship between toxoplasmosis and sex hormonal disturbances in seropositive males such as, testosterone (total, free) and Follicle Stimulating Hormone (FSH) levels.

Methods and Patients

Blood samples were collected from 248 apparently healthy males; their ages were between 21 and 60 years. They were attending the *In Vitro* Fertilization (IVF) center in Azadi Teaching hospital and the Central Blood Bank in Duhok city during the period from November 2016 to the end of July 2017. Before collecting the samples, a questionnaire was prepared and designed which covers various details. From each donor, 5ml of blood was withdrawn by vein puncture, placed in plain tubes and centrifuged. The serum was dispensed into 1ml labeled eppendorf tubes and stored at -20 °C until serological and hormonal tests were performed. Frequency percentage was

performed to determine the prevalence of affected persons by *T. gondii* and mean standard deviation was used to understand the reproductive male hormones concentrations. One-way ANOVA and chi-square tests were performed for statistical analyses. The independent t-tests and chi-square tests were performed to determine the association between toxoplasma and reproductive hormones and infertility[21-35].

Required Kits

- i. The Bioactiva Toxoplasma IgG ELISA (TOXG01) Kit was used.
- ii. The Bioactiva Toxoplasma IgM ELISA (TOXM02) Kit was used.
- iii. The Monobind Testosterone Enzyme Immunoassay Test Kit (3725-300) was used.
- iv. The Monobind Free Testosterone Enzyme Immunoassay Test Kit (5325-300) was used.
- v. The Monobind FSH Enzyme Immunoassay Test Kit (425-300) was used.

Results

The mean age of the studied participants was 37.29 ± 8.07 years. 22.98% of male samples appeared to be affected by *T. gondii* infection. The mean concentrations of the male reproductive hormones were 6.15 ± 1.03 ng/ml (total testosterone), 14.81 ± 4.71 pg/ml (Free Testosterone), 4.84 ± 2.20 IU/ml (Folliclestimulating hormone), 0.54 ± 0.22 (IgM), and 0.62 ± 0.75 (IgG). A significant difference was found in the samples in terms of infertility ($p < 0.0001$), while p-values for testosterone ($P > 0.05$), free testosterone ($P > 0.05$) and follicle-stimulating hormone ($P > 0.05$), were non-significant.

Conclusion

The present study showed a high rate of *T. gondii* among males in Duhok city with significant difference ($P < 0.05$) in infertility between *T. gondii* seropositive and seronegative samples(Tables 3-5).

Table 3: The seropositivity of *T. gondii* with respect to age and other factors.

Patients' Characteristics	Rate of Infection (N, %)				
	Overall Seropositive	22-30 Year	31-40 Year	41-50 Year	51-60 Year
Rate of infection	57(22.98)	14(5.6)	28(11.3)	14(5.6)	1(0.4)
Residence Urban Rural	26(10.5)	8(3.2)	12(4.8)	5(2.0)	1(0.4)
	31(12.5)	6(2.4)	16(6.4)	9(3.6)	0(0.0)
Occupation Public Private	29(11.6)	9(3.6)	15(6.0)	4(1.6)	1(0.4)
	28(11.3)	5(2.0)	13(5.2)	10(4.0)	0(0.0)
Fertility Fertile Infertile	53(21.3)	11(4.4)	27(10.8)	14(5.6)	1(0.4)
	4(1.6)	3(1.2)	1(0.4)	0(0.0)	0(0.0)
Blood groups					
O+	18(7.2)	7(2.8)	9(3.6)	2(0.8)	0(0.0)

O-	1(0.4)	1(0.4)	0(0.0)	0(0.0)	0(0.0)
A+	9(3.6)	2(0.8)	4(1.6)	2(0.8)	1(0.4)
A-	1(0.4)	0(0.0)	1(0.4)	0(0.0)	0(0.0)
B-	21(8.4)	3(1.2)	13(5.2)	5(2.0)	0(0.0)
B+	2(0.8)	0(0.0)	1(0.4)	1(0.4)	0(0.0)
AB+	4(1.6)	1(0.4)	0(0.0)	3(1.2)	0(0.0)
AB-	1(0.4)	0(0.0)	0(0.0)	1(0.4)	0(0.0)

Table 4: The mean concentration of TTH and FTH in the sera of the studied males infected with toxoplasmosis using ELISA according to the age group.

Age Group Year	IgM+			IgG+		
	N	TTH ng/ml	FTH pg/ml	N	TTH ng/ml	FTH pg/ml
22-30	1	9.3	30.5	12	7.68	21.44
31-40	6	6.09	17.39	22	6.72	17
41-50	8	6.04	13.71	7	8.38	24.66
51-60	1	5.76	13.65	0	0	0
Total	16			41		

Table 5: The mean concentration of TTH and FTH in the sera of the studied males infected with toxoplasmosis using ELISA according to the fertility.

Fertility	IgM+			IgG+		
	TTH ng/ml	FTH pg/ml	N	TTH ng/ml	FTH pg/ml	N
Fertile	6.1	13.72	13	7.27	19.5	39
Infertile	7.29	26.61	3	7.61	21.72	2
Total			16			41

Discussion

Seropositivity of toxoplasma gondii according to age




Figure 2: The distribution of studied males according to their seropositivity and seronegativity against toxoplasmosis using ELISA.

The current study showed out of 248 apparently healthy males from different ages (22-60 years) enrolled in this study, 57(22.98%) were seropositive for *T. gondii* antibodies. The rate of the infection increased with the increase in the age from 5.6% in the 22-30 age group to 11.3% in the 31-40 age group up to the age group 41-50 at which was decreased to 5.6%, then dramatically decreased to 0.4% at the age group 51-60. Various rates of seropositivity of *T. gondii* Abs using different serological tests have been reported among different age groups in previous

studies performed in different parts of Kurdistan and Iraq, in some of them higher rates were reported, while in others lower rates than that reported in this study were reported, this might be due to sample size, method of calculation, residency, age, type of test used, or other factors[36-43] (Figures 2-3).




Figure 3: The distribution of the studied males according to their seropositivity of anti *Toxoplasma* IgG and IgM antibodies using ELISA.

The distribution of toxoplasmosis using ELISA IGG and IGM according to blood group

This study showed that the overall seropositivity of toxoplasmosis among different blood groups was 22.98%.

Regarding blood groups, males of group B+aged 31-40 years showed the highest rate(5.2%) followed by 3.6% in males of blood group O+ of same ages.The present results partly agree with those ofwho found the highest percentage of seropositivity of *T. gondii* Abs in males of blood groups O+ and AB which were 35.8% and 38%, respectively[44-50].

Seropositivity of toxoplasma gondii according to residency

Regarding residency, the results of the current study on seropositivity of *T. gondii* in rural and urban areas, showed a high percentage in rural areas[51-53]. It is evident that people living in rural areas are more likely to be affected by different types of infectious microorganism such as, *T. gondii* due to more contact with animals. Regarding the age, ages from 31-40 and 41-50 years showed, higher rates of Toxoplasma Abs(6.4 and 3.6%, respectively) as compared to urban inhabitants[54-58](Figures 4A&4B).




Figure 4a: The distribution of the studied males according to seropositivity of anti-*Toxoplasma* IgG and age.




Figure 4b: The percentage distribution of the studied males according to seropositivity of anti-*Toxoplasma* IgM and age.

Seropositivity of toxoplasma gondii according to occupation

The results of the present study showed that toxoplasmosis seropositivity was different statistically with respect to occupation for difference of IgG and IgM for official and private occupation. The ages 31-40, 20-30, and 41-50 years in public sectors showed higher percentage of seropositivity (6.0%, 3.6%,

and 1.6%), respectively and the age group 31-40 years in private sector (5.2%)[59-65].

Effects of *T. gondii* on humans

The current study concentrated on the effects of *T. gondii* on male reproductive hormones including testosterone, free testosterone, an FSH, as well as, their impact on male fertility. The different aspects of scrutinized characteristics of the patients are analyzed in the following sections(Figures5-7).




Figure 5: The distribution of seropositivity of toxoplasmosis IgG and IgM Abs among blood groups in studied males.




Figure 6: The distribution of the studied males according to Seropositivity of IgG and IgM Antibodies among fertile and infertile males.




Figure 7: The distribution of the studied males according to seropositivity of toxoplasmosis and occupation.

The relation between *T. gondii* and reproductive hormones and fertility in males

The present study showed that there is a significant difference of male reproductive hormones between fertile and infertile males. Male infertility is caused by acquired or congenital urogenital abnormalities, infections in urogenital tract, increase in scrotum temperature, endocrine disturbances, hereditary abnormalities, and immunological factors [66]. Infectious agents such as bacteria, fungi, viruses and parasites are able to interfere with reproductive hormones and functions in male and female populations. The different organs of the male reproductive tract including testis, epididymis, or male accessory sex glands could be affected by infections[67-80](Tables 6-8).

Table 6: The relation between mean concentration of FSH in the sera of the studied males infected with toxoplasmosis from different ages using ELISA test.

Age Group	N	IgM+		N	IgG+	
		FSH (IU/ml)	FSH (IU/ml)		FSH (IU/ml)	FSH (IU/ml)
22-30	1	5.5		12	6.67	
31-40	6	5.07		22	4.92	
41-50	8	5.15		7	7.85	
51-60	1	4.33				
Total	16			41		

Table 7: The mean concentration of FSH in the sera of the studied males infected with Toxoplasmosis using ELISA according to the fertility.

Fertility	IgM+			IgG+		
	FSH (IU/ml)	N	%	FSH (IU/ml)	N	%
Fertile	4.68	13	81.25%	5.9	39	95.12%
Infertile	6.87	3	18.75%	6.54	2	4.88%
Total		16			41	

Table 8: The association of male reproductive hormones with *T. gondii* and fertility in men aged 22-60 years old.

Reproductive hormones		P-value (22-60) Years	
Testosterone *		P>0.05	
Free testosterone *		P>0.05	
follicle-stimulating hormone (FSH) *		P>0.05	
Fertility **		<0.0001	
Age groups**	Fertility N (%)		<0.0001
	Fertile	Infertile	
20-30	46 (73.0)	17 (27.0)	
31-40	98 (93.3)	7 (607)	
41-50	66 (98.5)	1 (1.5)	
51-60	12 (92.3)	1 (7.7)	

*ANOVA one way and **chi-square tests were performed for statistical analyses

The present study showed that the males infected with toxoplasmosis had the highest significant differences(P<0.001) with the mean concentration of serum FTH. Males with seropositive anti-Toxoplasma IgG antibodies and anti- Toxoplasma

IgM antibodies revealed the highest mean concentration levels of 19.61pg/ml and 16.13pg/ml for FTH hormones. While TTH and FSH hormone showed nonsignificant difference (P>0.05) for the mean concentration between acute and chronic males infected with toxoplasmosis[81-100] (Table 9).

Table 9: The association in men according to residency and occupation.

characteristics of Males	Fertile N (%)	Infertile N (%)	P-value
Residence	114(51.4)	11(42.3)	0.383*
	108(48.6)	15(57.76)	
Occupation	125(56.3)	18(69.2)	0.207*
	97(43.7)	8(30.8)	

*chi-square test was performed

Conclusion

From the present study, the followings are concluded:

- The overall seropositivity rate of *T. gondii* was 22.98% in the tested sample, 6.45% with acute and 16.35% with chronic toxoplasmosis.
- Regarding age, the highest rate (11.3%) of toxoplasmosis was among the age group 31-40years.
- Males with O+ and B+ blood groups showed higher percentage of infection with Toxoplasma in comparison to other blood groups[101-130].
- Infertile males had higher percentage of both acute and chronic toxoplasmosis(Figures 8-9).




Figure 8: The distribution of Toxoplasmosis IgG and IgM Abs seropositivity according to residency.

V. Official workers showed a higher rate (51.22%) of chronic toxoplasmosis than private worker (48.78%), while acute infection was higher (56.25%) in private workers than official (43.75%).

VI. Rural inhabitants revealed high rates of both acute and chronic toxoplasmosis, in comparison to urban inhabitants.




Figure 9: The mean concentration of sex hormones (TT, FT and FSH) in the sera of the studied males infected with acute (IgM) and chronic (IgG) toxoplasmosis using ELISA.

VII. The age group 22-30 year scored higher mean concentration of TT and FT in both acute and chronic infected males, while the age group 51-60 years showed a lower mean concentration of TT and FT in both acute and chronic infected males[131-140].

VIII. The TT and FT hormone levels recorded highest significant differences between fertile and infertile males. According to acute toxoplasmosis the infertile males showed a higher mean concentration of both TT and FT hormones. While, fertile cases, showed lowest significant differences.

IX. FSH showed higher value (6.87 and 6.54 IU/ml) and (4.68 and 5.9IU/ml) for acute and chronic toxoplasmosis in infertile and fertile men, respectively[141-152].

References

1. Al-Saadii, SHM (2013) The Effect of Toxoplasmosis on The Level of Some Male Sex Hormones in Samples from National Blood Transfusion Center/Baghdad M.Sc. thesis. College of Science/ University of Baghdad.
2. Bassad, Aboody Al (2015) Prevalence study of toxoplasmosis among males blood donors in thi-qar province-Iraq. IMPACT: International Journal of Research in Applied, Natural and Social Sciences (IMPACT: IJRNSS) 3(11): 73-82.
3. Al-Bajalan RR, Al-Nasiri FS, Mahmood SM (2015) Detection of Toxoplasma gondii by Latex and ELISA Test in Infertile and Fertile Men in Kalar City, Kurdistan Region, Iraq. International Journal of Current Microbiology and Applied Sciences 4(10): 570-585.
4. Abdoli A (2014) Toxoplasma, testosterone, and behavior manipulation: the role of parasite strain, host variations, and intensity of infection. Frontiers in biology 9(2): 151-160.
5. Abdoli A, Dalimi A, Movahedin M (2012) Impaired reproductive function of male rats infected with Toxoplasma gondii. Andrologia 44(s1): 679-687.
6. Abdul-Aziz AI, Zghair KH (2014) Study of Epidemiology of Toxoplasmosis in Hemodialysis Patients in Baghdad Hospitals. Iraqi Journal of Science 55(3B): 1236-1242.
7. Abdul-Lateef HI (2011) The levels of IFN-gamma, IL-12 And Testosterone Hormone in Persons with Asymptomatic Toxoplasmosis. M.Sc.Thesis. College of Science in Medical Microbiology, University of Baghdad: 102.
8. Abdulla HE, Al-Bashier NM, Al-Kawaz U, Al-Shuwaikh AMA, Abood AS (2015) Cross-sectional study of infertile males with Toxoplasmosis in Baghdad province. International Journal of Scientific and Engineering Research 6(1): 254-259.
9. Abu-Madi MA, Behnke JM, Dabritz HA (2010) Toxoplasma gondii seropositivity and co-infection with torch pathogen in high risk patients from Qatar. Am J Trop Med Hyg 82(4): 626-633.
10. Abubaker S, Dakhil V (2008) Seroprevalence of Toxoplasmosis in Women with Spontaneous Abortion. Zanco Journal of Medical Sciences 12(2): 48-54.
11. Ahmed SA, Gogal RM, Walsh JE, Saunders G (1996) Estrogen induces defects in T cell functions of mice. Fed Am Soc Exp Biol 10: 961.
12. Al-Dalawi NKE (2007) Hormonal disturbances in suddenly and previously aborted women affected with toxoplasmosis in Baghdad province. M.Sc. Technical Foundation College of Health and Medical Technology. Technical Foundation. PP. 125.
13. Al-Dujaily, KYA (1998) Seroepidemiological study of Toxoplasmosis among Aborted women in Baghdad. M.Sc., College of Veterinary Medicine/ University of Baghdad. PP. 123.
14. Jubori AL, AR M, (2005) Serological of toxoplasmosis in Kirkuk province. MSc Thesis. College of Health and Medical Technology. PP. 82.
15. Al-Kaysi AM, NM Ali (2010) Serological and biochemical study of HBV, HCV, HIV and toxoplasmosis infection among blood donors in Iraq. Egyptian Journal of Comparative Pathology And Clinical Pathology 23(1): 1-9.
16. Al-Qurashi AR (2004) Seroepidemiological study of toxoplasmosis in rural areas in the eastern region of Saudi Arabia. J Egypt Soc Parasitol 34(1): 23-34.
17. Al-Rawi KH (2009) Detection of B gene from blood of pregnant and abortive women infected with Toxoplasma gondii. Ph.D. Thesis. Collage of Science. University of Baghdad. PP. 125.
18. Al-Saadi AaM, Qasim MJ, Jebri NN (2013) Affliction with toxoplasmosis among infertile men in Kamal Al-Samaraee Hospital. Baghdad City. Al-Taqani 26(3): 56-63.
19. Al-Saeed ATM, Eassa SH, Murad MA (2016) Detection of toxoplasmosis among women with abortion using molecular and serological tests in Duhok City. DMJ 10(2): 56-68.
20. Al-Shikhly MA (2010) Early detection of toxoplasmosis percentage in premarital females by immunological methods. M.Sc. Thesis. College of Science. University of Baghdad pp. 147.
21. Al-Shikhly MA (2008) Serological study of Toxoplasma gondii antibodies in some Universities students in Baghdad province. M.Sc. thesis. College of Science. University of Baghdad pp. 102.
22. Al-Timimi RL (2004) Detection of toxoplasmosis among different groups of aborted women during gestational age of pregnancy. Diploma, Thesis. College of Health and Medical Technology pp. 77.
23. Al-Ubaydi GT (2004) Toxoplasmosis in pregnant women and its relation with some parameters. M.Sc. Thesis. College of Science. University of Mosul. pp. 127.
24. Al-Warid HS, Al-Qadhi BN (2012) Evaluation of progesterone and estrogen hormonal levels in pregnant women with toxoplasmosis. Eur J Sci Res 91(4): 515-519.
25. Altigany I, Ahmed SA, Abdrabo AEA (2014) Assessment of Testosterone Hormone Level in Patients with Toxoplasmosis Attending Omdurman Military Hospital. International Journal of Science and Research (IJSR) 4(12): 52-54.
26. Alwachi SN (2003) Biology. Daralfiker for printing and publishing, Amman, Jordan. (In Arabic). PP. 270.

27. Arantes TP, Lopes WD, Ferreira RM, Pieroni JS, Pinto VM, et al. (2009) Toxoplasma gondii: Evidence for the transmission by semen in dogs. *Experimental parasitology* 123(2): 190-194.
28. Arens J, Barnes K, Crowley N, Maartens G (2007) Treating AIDS-associated cerebral toxoplasmosis-pyrimethamine plus sulfadiazine compared with cotrimoxazole, and outcome with adjunctive glucocorticoids: South African Medical Journal 97(10): 956-958.
29. Ashraf M, Abdul-Haleem SA (2010) Seroprevalence of Anti-Toxoplasma gondii IgG and IgM Among Pregnant Women in Sana'a Capital and Capital Trusteeship. *Scientific Journal of King Faisal University (Basic and applied Sciences)* 11(2):179-188.
30. August JR (2009) Consultations in Feline Internal Medicine, Elsevier Health Sciences6: PP. 208.
31. Bahia-Oliveria LM, Jones JL, Azevedo-Silva J, Aleves CC, Orefice F, et al. (2003) Highly endemic, waterborne toxoplasmosis in north Rio de Janeiro state, Brazil. *Emerg Infect Dis* 9(1):55-62.
32. Bakre HM (2016) Serological tests and polymerase chain reaction for detection of Toxoplasma gondii infection in women attending for premarital examination. *Zanco Journal of Medical Sciences* 20(3): 1476 -1484.
33. Bakre HM, Hussain SK, Ali SK (2014) Toxoplasma gondii infection in patients with schizophrenia. *Zanco Journal of Medical Sciences* 19(1):874-879
34. Bassil N, Alkaade S, Morley JE (2009) The benefits and risks of testosterone replacement therapy: a review. *Ther Clin Risk Manag* 5(3): 427-448.
35. Bhopale GM (2003) Pathogenesis of toxoplasmosis. *Comparative immunology, microbiology and infectious diseases* 26(4): 213-222.
36. Bobić B, Klun I, Nikolić A, Djurković-Djaković O (2012) Toxoplasma gondii-infection in south-east Europe: epidemiology and epizootiology. *Toxoplasmosis-Recent Advances*: 37-54.
37. Brown WM, Hines M, Fane BA, Breedlove SM (2002) Masculinized finger length patterns in human males and females with congenital adrenal hyperplasia. *HormBehav* 42(4): 380-386.
38. Buffolano W, Gilbert RE, Holland FJ, Fratta D, Palumbo F, et al. (1996) Risk factors for recent Toxoplasma infection in pregnant women in Naples. *Epidemiol Infect* 116(3): 347-351.
39. Buxton D (1998) Protozoan infections (Toxoplasma gondii, Neospora caninum and *Sarcocystis* spp.) in sheep and goats: recent advances. *Veterinary research* 29(3): 289-310.
40. Carruthers VB (2002) Host cell invasion by the opportunistic pathogen Toxoplasma gondii. *Acta tropica* 81(2): 111-122.
41. Cartron JP (1994) Defining the Rh blood group antigens: biochemistry and molecular genetics. *Blood Rev* 8(4): 199-212.
42. Coêlho RA, Kobayashi M, Carvalho LB Jr (2003) Prevalence of IgG antibodies specific to Toxoplasma gondii among blood donors in Recife, Northeast Brazil. *Rev Inst Med Trop Sao Paulo* 45(4): 229-231.
43. Colosi HA, Jalali-Zadeh B, Colosi IA, Simon LM, Costache CA (2015) Influence of Toxoplasma gondii Infection on Male Fertility: A Pilot Study on Immunocompetent Human Volunteers. *Iran J Parasitol* 10(3): 402-409.
44. Cook AJ, Gilbert RE, Buffolano W (2000). Sources of Toxoplasma infection in pregnant women: European multicentre case-control studyCommentary: Congenital toxoplasmosis-further thought for food. *Bmj* 321(7254): 142-147.
45. Cox FE (2002) History of human parasitology. *Clin Microbiol Rev* 15(4): 595-612.
46. Craven AJ, Nixon AJ, Ashby MG, Ormandy CJ, Blazek K, et al. (2006) Prolactin delays hair regrowth in mice. *J Endocrinol* 191(2): 415-425.
47. Dalgıç N (2008) Congenital Toxoplasma gondii infection. *Marmara Medical Journal* 21(1): 89-101.
48. Dalimi A, Abdoli A (2013) Toxoplasma gondii and male reproduction impairment: a new aspect of toxoplasmosis research. *Jundishapur Journal of Microbiology* 6(8).
49. Drop SL, De Waal WJ, De Muinck Keizer-Schrama SM (1998) Sex Steroid Treatment of Constitutionally Tall Stature. *Endocrine Reviews* 19(5): 540-558.
50. Dubey J (2007) The history and life cycle of Toxoplasma gondii. *Toxoplasma gondii*: 1-17.
51. Dubey J (2009) History of the discovery of the life cycle of Toxoplasma gondii. *Int J Parasitol* 39(8): 877-882.
52. Dubey J, Hill D (2002) Toxoplasma gondii: Transmission and Prevention. *Clin Microbiol Rev* 8(10): 634-640.
53. Dubey J, Jones J (2008) Toxoplasma gondii infection in humans and animals in the United States. *Int J Parasitol* 38(11): 1257-1278.
54. Dubey JP (2008) The history of Toxoplasma gondii-the first 100 years. *J Eukaryot Microbiol* 55(6): 467-475.
55. Eslamirad Z, Hajhosseini R, Ghorbanzadeh B, Alimohammadi M, Mosayebi M, et al., (2013) Effects of Toxoplasma gondii Infection in Level of Serum Testosterone in Males with Chronic Toxoplasmosis. *Iran J Parasitol* 8(4): 622-626.
56. Eslamirad Z, Hajhosseini R, Gorbanzadeh B, AlmasiHashianee A (2014) Relationship between blood testosterone level and lipid profile among a group of men with and without Toxoplasma IgG antibody referred to pre-marital clinics in Arak city, 2012. *Arak Medical University Journal* 16(12): 1-8.
57. Flegr J (2007) Effects of Toxoplasma on human behavior. *Schizophr Bull* 33(3): 757-760.
58. Flegr J, Hrdá S, Kodym P (2005) Influence of latent/asymptomatic toxoplasmosis on body weight of pregnant women. *Folia Parasitol (Praha)* 52(3): 199-204.
59. Flegr J, Lindová J, Kodym P (2008) Sex-dependent toxoplasmosis-associated differences in testosterone concentration in humans. *Parasitology* 135(4): 427-431.
60. Garcia JL (2009) Vaccination concepts against Toxoplasma gondii. *Expert Rev Vaccines* 8(2): 215-225.
61. Garratty G, Postoway N, Nance SJ, Brunt DJ (1984) Spontaneous agglutination of red cells with a positive direct antiglobulin test in various media. *Transfusion* 24(3): 214-217.
62. Gazzinelli RT, Amichay D, Sharton-Kersten T, Grunwald E, Farber JM, et al. (1996) Role of macrophage-derived cytokines in the induction and regulation of cell mediated immunity to Toxoplasma gondii. *Current Topics in Microbiology and Immunology* 219: 127-139.
63. Giannoulis C, Zournatzi B, Giomisi A, Diza E, Tzafettas I (2008) Toxoplasmosis during pregnancy: a case report and review of the literature. *Hippokratia* 12(3):139-143.
64. Gill HS (1985) Occurrence of Toxoplasma gondii antibodies in Tanzanian blood donors. *East African medical journal* 62(8): 585-591.
65. Dohle GR, Colpi G, Hargreave T, Papp G, Jungwirth A, et al. (2005) EAU guidelines on male infertility. *European urology* 48(5): 703-711.
66. Gilot-Fromont E, Mercier A, Gotteland C, Richomme C, Aubert D, et al. (2012) The life cycle of Toxoplasma gondii in the natural environment, INTECH Open Access Publisher.

67. Guyton A (1986) Textbook of Medical Physiology (7th edn) Saunders. PA: 276, Philadelphia, USA.
68. Hakomori S (1999) Antigen structure and genetic basis of histoblood groups A, B and O: their changes associated with human cancer. *Biochim Biophys Acta* 1473(1): 247-266.
69. Hamad NR, Kadir MA (2013) Prevalence and comparison between the efficacy of different techniques for diagnosis of Toxoplasma gondii among women in Erbil province-Iraqi Kurdistan. *European Scientific Journal* 3(1): 901-908.
70. Henry L, Beverley KA (1976) Age and sex differences in the response of lymph node post-capillary venules in mice infected with Toxoplasma gondii. *British journal of experimental pathology* 57(3): 274-280.
71. Hill D, Dubey JP (2002) Toxoplasma gondii: transmission, diagnosis and prevention. *Clin Microbiol Infect* 8(10): 634-640.
72. Hill DE, Chirukandoth S, Dubey JP (2005) Biology and epidemiology of Toxoplasma gondii in man and animals. *Animal Health Res Rev* 6(1): 41-61.
73. Hodková H, Kolbeková P, Skallová A, Lindová J, Flegr J (2007) Higher perceived dominance in Toxoplasma infected men: the new evidence for role of increased level of testosterone in toxoplasmosis-associated changes in human behavior. *Neuro Endocrinol Lett* 28(2): 110-114.
74. Hunter CA, Sibley LD (2012) Modulation of innate immunity by Toxoplasma gondii virulence effectors. *Nat Rev Microbiol*; 10(11): 766-778.
75. Issit PD (1985) Applied Blood Group Serology (3rd edn). Montgomery Scientific, Miami, USA.
76. Jackson M, Hutchison W (1989) The prevalence and source of Toxoplasma infection in the environment. *Advances in parasitology* 28: 55-105.
77. Jafari R, Sadeghpour S, Sharifi F, Darani HY, Bagherpour B, et al. (2016) Anti-Toxoplasma IgM and IgG Seropositivity Among Individuals Referred to a Clinical Laboratory of Isfahan, Central Iran. *International Journal of Infection*; In press.
78. James WH (1996) Evidence that mammalian sex ratios at birth are partially controlled by parental hormone levels at the time of conception. *J Theor Biol* 180(4): 271-286.
79. James WH (1986) Hormonal control of sex ratio. *J Theor Biol* 118(4): 427-441.
80. Jassam FS (2010) Relationship between toxoplasmosis and testosterone hormone among schizophrenic patients in Baghdad. M. Sc. Thesis. College Council of Health and Medical Technology. PP 81.
81. Jones JL, Holland GN (2010) Annual burden of ocular toxoplasmosis in the United States. *Am J Trop Med Hyg* 82(3): 464-465.
82. Jones JL, Kruszon-Moran D, Wilson M, McQuillan G, Navin T, et al. (2001) Toxoplasma gondii infection in the United States: seroprevalence and risk factors. *Am J Epidemiol* 154(4): 357-365.
83. Jones JL, Muccioli C, Belfort R, Holland GN, Roberts JM, et al. (2006) Recently Acquired Toxoplasma gondii Infection, Brazil. *Emerging Infectious Disease journal-CDC* 12(4): 582-587.
84. Jones JL, Ogunmodede F, Scheftel J, Kirkland E, Lopez A, et al. (2003) Toxoplasmosis-related knowledge and practices among pregnant women in the United States. *Infectious diseases in obstetrics and gynecology* 11(3): 139-145.
85. Joseph B, Deepika S, Richard HS (2010) Toxoplasmosis. Web professional. Industry spotlight. Available.
86. Jumaian NF (2005) Seroprevalence and risk factors for Toxoplasma infection in pregnant women in Eastern Jordan. *Med Health J* 11(1-2): 45-51.
87. Jungwirth A, Diemer T, Dohle GR, Giwercman A, Kopa ZC, et al. (2015) Guidelines on Male Infertility (Report). Brussels.
88. Kadhim RA, Al-Awadi HM (2013) Changes in Testosterone, Progesterone and Prolactin Levels in Pregnant Women with Chronic Toxoplasmosis. *Med. J. Babylon* 10(3): 699-709.
89. Kaňková Š, Kodym P, Flegr J (2011) Direct evidence of Toxoplasma-induced changes in serum testosterone in mice. *Experimental parasitology* 128(3): 181-183.
90. Kapperud G, Jenum PA, Stray-pedersen B, melby KK, Eskild A (1996) Risk factor for Toxoplasma gondii infection in pregnancy. Results of a prospective case-control study in Norway. *Am. J. Epidemiol.* 144(4): 405-412.
91. Kareem S (2008) Prevalence, serodiagnosis and some immunological aspects of toxoplasmosis among women in Baghdad province. M.Sc. Thesis. College of health and Medical Technology. Technical Foundation pp: 104.
92. Kawashima T, khin-Sane W, Kawabata M, Barzaga N, Matsuda H, et al. (2000) Prevalance of antibodies to Toxoplasma gondii among urban and rural residents in the Philippines South east Asian. *J Trop Med Public Health* 31: 742-746.
93. Khalil HI (2008) Some aspects in Seroprevalence, diagnosis and influence of sex hormones on immunity during human toxoplasmosis. Ph.D. Thesis, College of Medicine. University of AL-Mustansiriyah pp. 173.
94. Kirkwood BR, Sterne JA (2010) Essential medical statistics, John Wiley and Sons.
95. Kittas C, Henry L (1980) Effect of sex hormones on the response of mice to infection with Toxoplasma gondii. *British journal of experimental pathology* 61(6): 590-600.
96. Lappalainen M, Hedman K (2004) Serodiagnosis of toxoplasmosis. The impact of measurement of IgG avidity. *Ann Ist Super Sanita* 40(1): 81-88.
97. Larosa DF, Stumhofer JS, Gelman AE, Rahman AH, Taylor DK, et al. (2008) T cell expression of MyD88 is required for resistance to Toxoplasma gondii. *Proceedings of the National Academy of Sciences* 105(10): 3855-3860.
98. Lecolier B, Grynberg H, Freund M (1990) Absence of relationship between Toxoplasma gondii antibodies and blood group in pregnant women in France. *European Journal of Clinical Microbiology and Infectious Diseases* 9(2): 152-153.
99. Leite M, Siciliano S, Rocha LSA, Justa TR, César KR, et al. (2008) Correlation between specific IgM levels and percentage IgG-class antibody avidity to Toxoplasma gondii. *Revista do Instituto de Medicina Tropical de São Paulo* 50(4): 237-242.
100. Liesenfeld O, Montoya JG, Tathineni NJ, Davis M, Brown BW, et al. (2001) Confirmatory serologic testing for acute toxoplasmosis and rate of induced abortions among women reported to have positive Toxoplasma immunoglobulin M antibody titers. *American journal of obstetrics and gynecology* 184(2): 140-145.
101. Liu Q, Wang ZD, Huang SY, Zhu XQ (2015) Diagnosis of toxoplasmosis and typing of Toxoplasma gondii. *Parasites and vectors*; 8(1): 292.
102. López R, Fano R, Contreras R, Font L (1993) Anticuerpos IgG anti-Toxoplasma gondii en Cubanos donantes de sangre. *Rev Lat Amer Microbiol* 35: 207-210.

103. Luft BJ, Chua A (2000) Central nervous system toxoplasmosis in HIV Pathogenesis, diagnosis, and therapy. *Curr Infect Dis Rep* 2(4): 358-362.
104. Lutchmaya S, Baron-Cohen S, Raggatt P, Knickmeyer R, Manning JT (2004) Second to fourth digit ratios, fetal testosterone and estradiol. *Early Human Dev* 77(1-2): 23-28.
105. Machattie C (1938) Notes on two cases of naturally occurring toxoplasmosis in Baghdad. *Tran. Roy Trop Hyg* 23:373-376.
106. Malarvizhi A, ViswanathanT, LavanyaV, Moorthy K (2012) Seroprevalence of Toxoplasma gondii in pregnant women. *Journal of public health and epidemiology* 4(6): 170-177.
107. MalikA, RizviM, Khan F, Khan N, Rabbani T, et al. (2014) Toxoplasma gondii in women with bad obstetric history and infertility: a five-year study. *Asian Pacific Journal of Tropical Disease* 4(1): S236-S239.
108. Markus MB(2003) Toxoplasma gondii. In: MiliotisMDand BIER JW(Eds.) International handbook of foodborne pathogens. New York: CRC Press, pp. 511-523.
109. Martinez-Garcia F, Regadera J, Mayer R, Sanchez S, Nistal M (1996) Protozoan infections in the male genital tract. *J urol* 156(2): 340-349.
110. Midtvedt T, Vaage L (1989) Relationship between Toxoplasma gondii antibodies and blood group. *Eur J Clin Microbiol Infect Dis* 8(6): 575-576.
111. Montoya JG, OLiesenfeld (2004) Toxoplasmosis. *The Lancet* 363(9425): 1965-1976.
112. Montoya JG, JSRemington (1995) Studies on the serodiagnosis of toxoplasmic lymphadenitis. *Clin infect dis* 20(4): 781-789.
113. Montoya JG, FRosso (2005) Diagnosis and management of toxoplasmosis. *Clin perinatal* 32(3): 705-726.
114. Mooradian AD, Morley JE, Korenman SG (1987) Biological actions of androgens. *Endocr Rev* 8(1): 1-28.
115. Neave N, LaingS, Fink B, Manning JT(2003) Second to fourth digit ratio, testosterone and perceived male dominance. *Proc Biol Sci* 270(1529): 2167-2172.
116. Nelson RJ (2005) An introduction to behavioral endocrinology (3rdedn). Sunderland, Mass: Sinauer Associates.Inc. pp. 143.
117. Niazi AD, MW N, saif SA, AbbassAF, Gzar(1992)Prevalence of Toxoplasma antibodies in Iraqi population. *J Fac Med Baghdad* 34(3): 255-261.
118. Ökten A, Kalyoncu M, YarışN(2002) The ratio of second-and fourth-digit lengths and congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Early hum dev* 70(1): 47-54.
119. OktenliC, L Dogansit,Ozgurtas(2004) Transient hypogon- adrotrophic hypogonadism in males with acute toxoplasmosis: suppressive effect of interleukin-1 β on the secretion of GnRH. *Human Repro* 19(4):859-866.
120. Pappas G, Roussos N, Falagas ME(2009) Toxoplasmosis snapshots: global status of Toxoplasma gondii seroprevalence and implications for pregnancy and congenital toxoplasmosis. *Int J Parasitol* 39(12): 1385-1394.
121. Pellati D, Mylonakis I, Bertoloni G, Fiore C, Andrisani A et al. (2008) Genital tract infections and infertility. *Eur J Obstet Gynecol Reprod Biol* 140(1): 3-11.
122. Pergola G, Cascone A, Russo M (2010) Acute pericarditis and myocarditis by Toxoplasma gondii in an immunocompetent young man: a case report. *Infez Med* 18(1): 48-52.
123. Petersen E (2007) Toxoplasmosis. *Seminars in Fetal and Neonatal Medicine* 12(3): 214-223.
124. Race RR, R Sanger (1975) Blood groups in man. (6thedn). Oxford, Published by Blackwell Scientific Publications.
125. RäisänenS (1978) Toxoplasmosis transmitted by blood transfusions. *Transfusion* 18(3): 329-332.
126. Razzaq A H, SAWais, AYSaeid(2005)Toxoplasmosis: the innocent suspect of pregnancy wastage in Duhok, Iraq. *East Mediterr health J* 11(4): 625- 632.
127. Remington JS, EsmontsGD (1990)Toxoplasmosis In: JS Remington and JOKlein(Eds). *Infectious Disease of the fetus and New born Infant*.WB Saunders. London, pp. 89-195.
128. Remington JS, RMcLeod, PThulliez, GDesmonts(2005)Toxoplasmosis. In: JS Remington. *Infectious Diseases of the fetus and newborn infant*,WBSaunders Company Philadelphia, pp.947-1091.
129. Rodrigues A, SUezatos, MVono, TPandossio, LSpiegiorinA, OlianiD, et al. (2011)Association between anti- Toxoplasma gondii antibodies and ABO blood group system. *JVenom Anim Toxin Trop Dis* 17(2):184-189.
130. Roitt I, J Brostoff,D Male(2001)Parasitology and Vector biolgy, (2nd ed). Academic Press. pp165-178.
131. Sabin AB, Feldman HA (1948) Dyes as microchemical indicators of a new immunity phenomenon affecting a protozoon parasite (Toxoplasma). *Science* 108(2815): 660-663.
132. SalibayCC, CADung,FG Claveria(2008)Serological survey of Toxoplasma gondii infection among urban.(Manila)and suburban (Dasmarinas,cavite) Residents, Philippines *JprotozolRes* 18:26-33.
133. SenthamaraiS,SivasankariS,Apurba S, Sandhya B,KumudavathiM, et al. (2013)Seroprevalence of toxoplasmosis in pregnant women with bad obstetric history in a tertiary care hospital, Kanchipuram-a pilot study. *Disease* 3(9): 29-32.
134. SherA (1998) Ignition of the type 1 response to intracellular infection by dendritic cell-derived interleukin-12. *Eur cytokine netw* 9(3 Suppl): 65-68.
135. Shin DW, Cha DY, Hua QJ, Cha GH, Lee YH (2009) Seroprevalence of Toxoplasma gondii infection and characteristics of seropositive patients in general hospitals in Daejeon, Korea.*Korean j parasitology* 47(2): 125-130.
136. Siddiqui N, Shujatullah F, Khan HM, Rabbani T, Khan PA (2014) IgG avidity antibodies against Toxoplasma gondii in high risk females of reproductive age group in India.*Korean J parasitol* 52(5): 487.
137. Sinisi AA, Di Finizio B, Pasquali D, Scurini C, D Apuzzo A, et al.(2003) Prevalence of antisperm antibodies by SpermMARtest in subjects undergoing a routine sperm analysis for infertility. *Int J Androl* 16(5): 311-314.
138. Sukthana Y (2006) Toxoplasmosis: beyond animals to humans. *Trends Parasitol* 22(3): 137-142.
139. Swai ES, Schoonman L(2009) Seroprevalence of Toxoplasma gondii infection amongst residents of Tanga district in north-east Tanzania. *Tanzanhealth Res* 11(4): 205-209.
140. Szekeres-BarthoJ, Polgar B (2010)Chapter 8: Progesterone, Pregnancy, and Innate Immunity. In: KLEIN SL and Roberts, CW(eds.) *Sex hormones and immunity to infection*. Springer, USA.
141. Tenter AM, Heckerth AR, Weiss LM (2000) Toxoplasma gondii: from animals to humans. *Int j parasitol* 30(12): 1217-1258.
142. Terpsidis KI, Papazahariadou MG, Taitzoglou IA, Papaioannou NG, Georgiadis MP, et al. (2009) Toxoplasma gondii: reproductive

- parameters in experimentally infected male rats. *Expparasitol* 121(3): 238-241.
143. Tuck S, Francis R (2009) Testosterone, bone and osteoporosis. Advances in the management of testosterone deficiency. Karger 137: 123-132.
144. Uniprot(2017)Taxonomy-Piroplasmida(Piroplasmids).
145. Webster JP (2001) Rats, cats, people and parasites: the impact of latent toxoplasmosis on behaviour. *Microbesinfect* 3(12): 1037-1045.
146. Widmann FK(1990) Technical Manual. Washington DC: American Association of Blood Banks.
147. Wilson M, Remington JS, Clavet C, Varney G, Press C, et al.(1997) Evaluation of six commercial kits for detection of human immunoglobulin M antibodies to *Toxoplasma gondii*. The FDA Toxoplasmosis Ad Hoc Working Group. *J Clin Microbiol* 35(12): 3112-3115.
148. Wolf A, Cowen D, Paige BH (1939)Toxoplasmic encephalomyelitis: III. A new case of granulomatous encephalomyelitis due to a protozoan. *Am jpathol*15(6): 657-694.
149. Zaini RG, Ismail KA, Dahlawi H (2016) Seroprevalence of *Toxoplasma gondii* among AIDS Patients in Saudi Arabia. *World Journal of AIDS* 6(03): 81.
150. Zhiburt EB, Ionova AI, Danil'chenko VV, Serebrianaia NB, Bel'gesov NV, et al. (1996) The spread of antibodies to cytomegalovirus and *Toxoplasma* among donors of blood components. *ZhMikrobiol Epidemiol Immunobiol*1: 59-61.
151. Zhou P, Zhang H, Lin RQ, Zhang DL, Song HQ (2009) Genetic characterization of *Toxoplasma gondii* isolates from China. *InJParasitol*58(2):193-195.
152. Zuber P, Jacquier P (1994) Epidemiology of toxoplasmosis: worldwide status. *Schweiz Med Wochenschr Suppl* 65: (19S-22S).



This work is licensed under Creative Commons Attribution 4.0 Licens
DOI: [10.19080/AIBM.2018.11.555823](https://doi.org/10.19080/AIBM.2018.11.555823)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>