



Research Article

Volume 9 Issue 1 - July 2017
DOI: 10.19080/ARTOAJ.2017.09.555752

Agri Res & Tech: Open Access J

Copyright © All rights are reserved by Jayanth Kumar

Effect of Organic Priming on Germination and Vigour of Cotton (*Gossypium hirsutum* L.) Seed



Jayanth Kumar P*, Chaurasia AK and Bineetha M Bara

Department of Genetics and Plant breeding, Sam Higginbottom University of Agriculture Technology and Sciences, India

Submission: June 13, 2017; Published: July 28, 2017

*Corresponding author : Jayanth Kumar, Department of Genetics and Plant breeding, Sam Higginbottom University of Agriculture Technology and Sciences, India, Email: kumarjayanth86@gmail.com

Abstract

The experiment was conducted in Seed science post Graduate Laboratory, Department of Genetics and Plant Breeding, SHUATS, Allahabad, U.P. In order to standardize the best method of Organic priming specific to cotton, two methods of priming viz., Organic priming and hydropriming and they were evaluated by screening a range of durations and concentrations viz., T₀-Unprimed Control, T₁-Distilled water hydration (for 12 hrs), T₂-Cow urine at 2%, T₃-Cow urine at 6%, T₄-Cow urine at 10%, T₅-Curry leaf extract 6%, T₆-Curry leaf extract 10%. It found that all the Organic priming methods showed significance difference with the control and the highest germination %, seedling length (cm), seedling fresh weight (g), seedling dry weight (g) and vigour index were observed in T₃ for Cow urine at 6%. The study helps to improve the quality of seeds with the help of seed organic priming treatments which are cost effective and economic, nontoxic, ecofriendly sources.

Keywords: (*Gossypium hirsutum* L.); Organic priming; Cow urine; Curry leaf extract; Hydropriming

Introduction

Cotton (*Gossypium sp.*) the “white gold” and “king of fibers”, is closely linked to human civilization itself. The English term cotton derives its name from Arabic word “Quotri”, Dutch ‘Katoem’ and French ‘Cotton’. Archaeologists have discovered cotton fibers more than 4,000 years ago in coastal Peru and at Mohenjo-Daro in the Indus valley (Pakistan). But the origin of

cotton is shrouded in mystery. India is considered the home of arborium cottons. But *Gossypium* was possibly introduced into Western India from Arabia, Persia and Baluchistan. Cotton is cultivated in tropical and subtropical regions of more than seventy countries across the world and enjoys a predominant position amongst all cash crops in India.

Table 1: Analysis of variance for 8 seedling characters in cotton.

S.No.	Characters	Mean Sum of Squares		
		Replication (d. f. = 03)	Treatments (df=6)	Error (df=21)
1.	Germination percentage	6.000	78.643**	20.905
2.	Speed of germination	4.952	37.536**	11.833
3.	Root length	5.457	12.921**	1.390
4.	Shoot length	0.838	19.739**	1.265
5.	Seedling length	6.734	20.879**	2.022
6.	Seedling fresh weight	0.186	1.764**	0.175
7.	Seedling dry weight	0.004	0.198**	0.002
8.	Seed vigour index length	464.972	1798.270**	209.675
9.	Seed vigour index mass	30.372	2030.699**	36.332

**significant at 5% and 1% level of significance, respectively.




Figure 1: Germination percentages as Influence by different Organic priming treatments on cotton seeds.




Figure 2: Speed of germination as Influence by different Organic priming treatments on cotton seeds.




Figure 3: Root length (cm) as Influence by different Organic priming treatments on cotton seeds.

Due to environmental concerns, there is an urgent need to reduce the use of chemical fertilizers and pesticides in agriculture and horticulture and alternative to chemicals are being sought to improve crop establishment and health. One option is the use of organics nutrients or growth regulators to seed or roots, which may promote plant growth or provide diseases control through a variety of mechanisms, including supply of organic nutrients production of plant hormones, antibiotic or enzyme; induced systemic resistance; direct parasitism of plant pathogen or deleterious micro-organisms; or competition with pathogen for or nutrients. Further, organic seed is a crucial link in the chain from research to organic seed production and ultimate supply of high quality seed at reasonable price to the commercial seed producing farmers for promotion of organic seed production. Hence, the safe and feasible approach is the priming of seeds with organics which are safe, ecofriendly, economical and easily available. Organic seed priming provides hardiness to

high temperature, low moisture especially in semiarid tropics. It promotes faster germination, higher seedling vigour leading to higher crop productivity. The main benefits of organic seed treatments include increased phosphate levels, nitrogen fixation and root development (Figure 1-5 & Table 1).




Figure 4: Shoot length (cm) as Influence by different Organic priming treatments on cotton seeds.




Figure 5: Seedling length (cm) as Influence by different Organic priming treatments on cotton seeds.

Cow urine contains about 1.0% nitrogen, traces of P2O5 and 1.0% of K2O. Approximately 2400 to 2500L of urine are produced per year per animal. If this urine were not conserved, nitrogen in the urine, which is mainly in the form of urea, would be quickly lost as ammonia. It is also considered as a natural disinfectant and pest repellent and forms the main component of Panchagavya (an organic crop booster prepared and sprayed by Indian farmers) [1]. Organic seeds priming is more affordable so even small scale farmers can practice.

Curry leaves have been used for centuries almost in all the parts of the world. This herb has several medicinal properties. For instance, its leaves and bark can be used as a tonic, stomachic, stimulant and carminative. It can also help in reducing blood sugar if these leaves are consumed early in the morning in empty stomach. Curry leaves are also a good source of vitamin A and they provide a rich source of calcium. Curry leaves are a great source of various vitamins and minerals. These include vitamin C, Vitamin A, folic acid, niacin, thiamin and riboflavin. Each of these vitamins plays an important role in development. Vitamin C is important for strengthening the immune system. Thiamin is known to have a role in organ and nervous system development. Other than vitamins and minerals, studies have shown that curry leaves contain antioxidants which are useful against free radical damage and oxidative stress in the body.

Materials and Methods

The experiment was conducted in Seed science post Graduate Laboratory, Department of Genetics and Plant Breeding, SHUATS, Allahabad, U.P.

Genetically pure seeds of cotton hybrid JK CH 8665 BG 2 used for the study. The seeds were treated with cow urine at the concentration of 2, 6 and 10% and curry leaf extract at the concentration of 6 and 10% along with distilled water and dry seed as control. The seeds were tested for the standard germination test adopting between paper (BP) method and

sand method as per the ISTA rules. Locally available cow urine is used for organic seed priming which acts as growth promoter by preventing plant disease. 2, 6, and 10ml of cow urine were added separately in 100ml water to get 2%, 6%, 10% of solution which were used for seed treatment as per the required weight by volume ratio of seed to solution. Curry leaf extract solution was prepared by according to Sathish Chand Saini [2] and distilled water solution was prepared by according to Pill and Necker (2001). Seed to solution ratio of 1:0.5, were made and soaked to 8 hours for cotton. Then seeds were dried overnight (Table 2).




Table 2: Mean performance of cotton for 9 seedling character.

S.No	Treatments	Germination %	Speed of Germination	Root Length (cm)	Shoot Length (cm)	Sending Length (cm)	Fresh Weight of Sending (g)	Dry Vigour Index Length	Seed Vigour Index Length	Seed Vigour Index Mass
1	T0	70.25	69	11.15	10.63	22.66	5.47	1.11	165.38	77.46
2	T1	71.25	72	12.6	11.83	23.53	5.67	1.11	166.33	78.97
3	T2	71.5	70	12.55	11.51	25.34	6.64	1.15	180.76	81.97
4	T3	83.25	78	16.54	17.39	29.38	7.46	1.66	223.11	137.72
5	T4	72	72	12.34	11.63	23.34	6.39	1.13	162.82	81.3
6	T5	73.25	74	14.67	12.85	26.2	6.68	1.5	192.21	109.56
7	T6	72.5	70	12.62	12.81	24.57	6.4	1.16	177.98	84.32
G.MEAN		73.42	72.14	13.21	12.66	25.01	6.38	1.26	182.22	93.042
C.D. (5%)		6.723	5.058	1.733	1.654	2.091	0.615	0.065	21.293	80864
SE(m)		2.286	1.72	0.589	0.562	0.711	0.209	0.022	7.24	3.014
C.V.		6.227	4.759	8.925	8.881	5.688	6.553	3.502	7.99	6.478

The data collected from the experiments were analyzed statistically by the procedure prescribed by. The investigations are details of materials and using of experimental design in complete randomized block methods. Critical difference were calculated at 1% level wherever 'F' test was significant. The data on percentage of germination and seed infection were transferred into arc sine square root percentage values and the transferred data were used for statistical analysis. Absolute control treatment was compared with rest of the treatment by following ANOVA statistical analysis [3-10].

Result and Discussion

Germination (%)



A range of 70.25 to 83.25 percent was observed for germination percentage. The mean value for this parameter was 73.43 percent. Maximum germination percentage (83.25) was recorded with T₃-Cow urine at 6%. Whereas minimum significantly germination percentage followed by (73.25) with T₅-application of Curry leaf extract 6%. Minimum germination percentage was recorded by T₀ (70.25) with control (Figure 6-9).




Figure 9: Seed vigour index mass Influence by different Organic priming treatments on cotton seeds.

Speed of germination

A range of 69 to 78 percent was observed for germination percentage. The mean value for this parameter was 72.14 percent. Maximum germination percentage was recorded with T₃ Cow urine at 6% (78.00). Whereas minimum significantly germination percentage followed by T₅ with application of Curry leaf extract 6% (74.00). Minimum germination percentage was recorded by T₀ (69) with control.

Root length

The mean performance of seedling root length ranged from 11.15cm to 16.54cm with mean value of 13.21cm. Maximum root length (16.54cm) was recorded by T₃ with application Cow urine at 6% and it was followed by T₅ (14.67cm) with application of Curry leaf extract 6%. Minimum root length was recorded by T₀ (11.15cm) with control.

Shoot length

Among shoot length there exists a significant variation as influenced by organic priming methods and seeds primed with T₃ with application of Cow urine at 6% (17.39cm) has highest root length, control (10.63cm) has lowest root length. Seeds primed with cow urine at 2% and hydro primed seeds (11.83), curry leaf extract 10% (12.81) have statistically on par results [10-15].

Seedling length

The mean performance of seedling length ranged from 22.66cm to 29.38cm with mean value of 25.01cm. Maximum seedling length (29.38cm) was recorded by T₃ with application of Cow urine 6% and it was followed by T₅ (26.20cm) with application of Curry leaf extract 6%. Shortest seedling length was recorded by T₀ (22.66cm) with control [15-17].

Seedling fresh weight

The mean performance of seedling fresh weight ranged from 5.47g to 4.27g with mean value of 6.38g. Maximum seedling fresh weight (7.46g) was recorded by T₃ with application of Cow urine 6% and it was followed by T₅ (6.68g) with application of Curry leaf extract 6%. Lowest value of seedling fresh weight was recorded by T₀ (5.47g) with control.

Seedling dry weight

The mean performance of seedling dry weight ranged from 1.11mg to 1.66mg with mean value of 1.26mg. Maximum seedling fresh weight (1.66mg) was recorded by T₃ with application of Cow urine 6% and it was followed by T₅ (1.50mg) with application of Curry leaf extract 6%. Lowest value of seedling fresh weight was recorded by T₀ (1.11g) with control.

vigour index I

The mean performance of seedling vigour index length ranged from 165.38 to 223.11 with mean value of 181.22. Maximum seedling vigour index length (223.11) was recorded by T₃ with application of Cow urine at 6% and it was followed by T₅ (192.21) with application of Curry leaf extract 6%. Minimum seedling vigour index length was recorded by T₀ (165.38) with control.

Seed vigour index II

The mean performance of seedling vigour index mass ranged from 77.46 to 137.72 with mean value of 93.042. Maximum seedling vigour index mass (137.72) was recorded by T₃ with application of Cow urine at 6% and it was followed by T₅ (109.56) with application of Curry leaf extract 6%. Minimum seedling vigour index mass was recorded by T₀ (77.46) with control.

Conclusion

All the priming methods have positive influence on seed quality parameters of cotton individually but the effect of priming method was found significant. Speed of germination and Germination percentage (78 and 83.25%) respectively and were highest in Cow urine at 6% seeds and it was significantly low in unprimed (control) seeds (69 and 70.25%). However seedling attributes were also positively influenced by organic priming and highest seedling length (29.38cm) and seedling dry weight (1.66mg) was observed in Cow urine at 6% seeds followed by Curry leaf extract 6%. Seeds having seed vigour index length and seed vigour index mass (191.60 and 137.72mg) respectively and found to be lowest in unprimed seeds having seedling attributes (145.06 and 77.46mg respectively).

Organic Priming increases the germinability and vigour of cotton seeds, significantly in lab condition. Cow urine at 6% and Curry leaf extract 6% significantly increased the germination and vigour of cotton. Organic priming of the cotton seeds for 12hrs. Enhanced germinability, vigour of cotton seeds. These conclusions are based on the results of three months investigation and therefore further investigation is needed to

arrive at valid recommendations.

References

1. Tharmaraj K, Ganesh P, Kumar RS, Anandan A, Lolanjinathan K (2011) A critical review on Panchagavya-a boon to plant growth. International Journal of Pharma Biol Arch 2(6): PP1611-1614.
2. Saini SC, GBS Reddy (2013) Assessment of Quality of Curry Leaves (Murraya koenigii). International Journal of Pharmaceutical Science Invention 2(10): PP13-17.
3. Abdul-Baki AA, Anderson JD (1973) Vigour deterioration of soybean seeds by multiple criteria. Crop Science journal 13: PP630-633.
4. Arif M, Khan MTJ, Marwat B, Khan MA, (2008) Seed Priming Improves Emergence and Yield of Soybean. African Journal of Biotechnology 40(3): PP1169-1177.
5. Amarnath BH, Chaurasia AK, Kumar A, Chaurasia N, Vivekanad V, et al. (2015) Effect of priming with botanicals and animal waste on germination and seedling vigour in sorghum (*Sorghum bicolor* L.) seeds. Advances in Applied Science Research 6(10): PP73-77.
6. Ambika S, Balakrishnan K (2015) Enhancing germination and seedling vigour in cluster bean by organic priming. Academic Journal 10(8): 298-301.
7. Ambika S, Balakrishnan K, Sujatha K (2014) Enhancing the Seed Germination and Vigour in Coarse Cereals by Bovine Urines. Journal of Agroecology and Natural Resource Management 1(2): 40-43.
8. Bradford KJ, Steiner JJ, Trawatha SE (2002) Seed priming influence on germination and emergence of pepper seed lots. Crop Science 30: PP718-721.
9. Disegha GC, Izionworu VO (2014) Antifungal Activities of Curry Leaf (Murraya Koengii) Extract on Some Selected Fungi. Chemistry and Materials Research 6(11).
10. Jain P, Sharma RC, Bhattacharya P (2014) Effect of new organic supplement (Panchgavya) on seed germination and soil quality 186(4): 1999-2011.
11. Nawaz J, Hussain M, Jabbar A, Nadeem GA, Sajid M, et al. (2013) Seed Priming A Technique. International Journal of Agriculture and Crop Sciences.
12. Jayashri V, Karmore, Tomar GS (2015) Effects of seed priming methods on germination and seedling development of winter maize (*Zea mays* L.). Advance research journal of crop improvement 6(2): 88-93.
13. Kumar M (2014) Influence of Seed Priming with Urine, Phosphorus and Zinc on Maize (*Zea mays* L.) Yield in an Acid Soil of Northeast India. Indian Journal of Hill Farming 27(1): 78-80.
14. Mehta DK, Kanwar HS, Thakur AK, Thakur KS (2010) Influence of organicseed priming on germination and seedling quality in bell pepper (*Capsicum annum* L.). Journal of Hill Agriculture 1(1): 85-87.
15. Shankaranarayan R, Vijayakumar M, Rangaswamy P (1994) Cow urine for ideal seed germination in tamarind. India Hort 38(4): PP15.
16. Sutteesh VK, Jijeesh CM, Divya TP (2016) Evaluation Of Organic And Inorganic Pretreatments For Better Seed Germination and Seedling Vigour In Santalum album L.. Plant Archives 16(1): PP 143-150.
17. Uchida Y, Clough TJ, Kelliher FM, Hunt JE, Sherlock RR (2011) Effects of bovine urine, plants and temperature on N₂O and CO₂ emissions from a sub-tropical soil. Plant Soil 345: 171-186.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/ARTOAJ.2017.09.555752](https://doi.org/10.19080/ARTOAJ.2017.09.555752)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>