



Review Article

Volume 9 Issue 3 - August 2017
DOI: 10.19080/ARTOAJ.2017.09.555765

Agri Res & Tech: Open Access J
Copyright © All rights are reserved by : Imran

Traditional Rice Farming Accelerate CH₄ & N₂O Emissions Functioning as a Stronger Contributors of Climate Change



Imran^{1,2*}, Abdul Bari³, Roshan Ali³, Naeem Ahmad³, Zulfiqar Ahmad², Muhammad Idrees Khattak⁴, Akhtar Ali³, Fayaz Ahmad³, Inayatullah Khan¹ and Shahida Naveed⁵

¹Department of Agronomy, The University of Agriculture, Pakistan

²Climate Change Centre (CCC), The University of Agriculture, Pakistan

³Agriculture Research Institute Mingora Swat, Pakistan

⁴Department of Rural sociology, The University of Agriculture, Pakistan

⁵Associate professor of Botany, Govt. Girls degree college Karak, Pakistan

Submission: July 21, 2016; Published: August 10, 2017

*Corresponding author : Imran, Department of Agronomy, the University of Agriculture, Pakistan, Email: imranagrarian@aup.edu.pk

Abstract

Traditional methods of rice farming, like those practiced in the South Asian countries included Pakistan, contribute to greater GHG emissions than in other regions. These methods include constant saturation of rice fields (which depletes oxygen in the system), the addition of fertilizers (which provides nitrates that stimulate denitrification in bacteria), and increasing the acreage of rice fields which increases the area available for GHG emissions. Alternate methods such as modified irrigation and more efficient fertilizer use have proven to be effective in reducing GHG emissions. Another improved method of rice farming, Direct Seeded Rice (DSR) is also a step toward climate change and GHG mitigation strategy to be adopted and disseminated in the farming community of the rice growing countries. However it is a challenge to identify and improve ways of increasing rice yields while simultaneously reducing GHG emissions to widespread implementation of GHG-reducing practices due to limited outreach and awareness regarding the incentives and benefits of alternate methods of rice farming.

Keywords : Rice (*Oryza sativa L.*); Climate change; Greenhouse gases; Methane; Nitrous oxide

Introduction

Oryza sativa L. is an important kharif crop of Pakistan and ranked second to wheat as a staple food. Rice is occupying a prime position in the agricultural economy of Pakistan. In Asia, more than 3.5 billion people prefer rice for food and this demand for rice will be increased up to 70 percent in 2025 [1]. In Pakistan the area under rice cultivation is 2.25 million hectares, thus becoming the country's third largest crop in terms of area. Rice cultivation has been playing an effective role in uplifting the economic conditions of the common farmer community of the country. In Khyber Pakhtunkhwa rice is the second preferable cultivable crop to wheat and maize, grown under two different agro climatic conditions, i.e., plains and upper mountainous valleys. Out of the total rice cultivated area (64719ha), 81% comprised of cooler, high altitude areas of Malakand, Hazara and adjacent tribal

areas of Khyber Pakhtunkhwa [2]. The Intergovernmental Panel on Climate Change (IPCC) has confirmed that anthropogenic emissions of greenhouse gases (GHGs), which are those emitted by human activity, contribute to climate change and are at their highest levels in history. GHGs contribute to the warming of the planet by trapping radiation and releasing it near the surface of the earth. Agricultural practices emit over 50% of global non-carbon dioxide GHG emissions. About 7% of these emissions are a result of rice farming, and over 80% of GHG emissions from rice farming are produced in South and Southeast Asia (India, China, Indonesia, Thailand, Vietnam, and Burma). The warming of the planet, combined with the growing population expected to exceed 9 billion by 2050, threaten advancements made to achieve global food security and environmental sustainability.

The earth's surface, like all planetary bodies, releases infrared radiation. While this radiation usually travels through the atmosphere and escapes the surface, greenhouse gases (GHGs) that are present in the atmosphere, such as carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O) and even water (H₂O), can capture this radiation and release it back to the surface of the earth [3]. GHGs are emitted as a byproduct of many activities such as burning fossil fuels, generating electricity, and agriculture. While CO₂ is often the first one that comes to mind, CH₄ and N₂O are much stronger contributors of GHGs emissions. Rice farming is a significant source of human-produced non-CO₂ emissions. Because rice is grown in flooded systems, the soil remains waterlogged, preventing oxygen from entering the pores. Additionally, the decomposition of organic plant matter in the fields consumes and therefore, limits the amount of oxygen present. With limited oxygen in the soil, bacteria that can conduct anaerobic respiration, or "breathing" in the absence of oxygen, thrive. Two of the most common methods of such respiration in flooded rice fields are methanogenesis, the production of CH₄, and denitrification, the conversion of nitrates to nitrogen gas with N₂O as a byproduct. Paddy rice fields are particularly conducive to these processes not only because the soil is waterlogged and therefore lacks oxygen; but also because anaerobic bacteria rely on plant matter from rice for respiration.

Methanogenesis

Rice is grown in flooded systems, the soil remains waterlogged, preventing oxygen from entering the pores. Additionally, the decomposition of organic plant matter in the fields consumes and therefore, limits the amount of oxygen present. With limited oxygen in the soil, bacteria that can conduct anaerobic respiration, or "breathing" in the absence of oxygen, thrive. Two of the most common methods of such respiration in flooded rice fields are methanogenesis, and we call that the production or emission of methane gas (CH₄) from rice field [3]. Methane in rice paddies is produced by microscopic organisms that respire CO₂, like humans respire oxygen. More CO₂ in the atmosphere makes rice plants grow faster, and the extra plant growth supplies soil microorganisms with extra energy, pumping up their metabolism. Increasing CO₂ levels will also boost rice yields, but to a smaller extent than CH₄ emissions. As a result, the amount of CH₄ emitted per kilogram of rice yield will increase [4].

Denitrification

The conversion of nitrates to nitrogen gas with N₂O as a byproduct. Simply stated, when sufficient oxygen is not available in soil. Some microorganisms use oxygen from the nitrate ion (NO₃⁻). When oxygen is removed from nitrate, it leaves N gas, which can then be lost to the atmosphere. This is denitrification. Soil conditions that lead to denitrification typically occur when moisture is high or there is standing water, and conditions are warm enough to encourage microbial activity. Loss due to denitrification can be significant, especially in a wet year. When

the soil microorganisms need oxygen, they'll take it from the air or from nitrate if the water in the soil prevents them from getting it from the air. The soil does not have to be completely saturated for denitrification to occur in this way (Figure 1).




Figure 1: Graph showing GHG emission percentage, Source: Mitigation related agreements, Koyoto Protocol, Paris agreement.

Denitrification chemistry and occurrence

Denitrification is a microbial process in the soil that reduces N-containing compounds into their simplest forms. Nitrification makes N available to the plant. Denitrification is one of the primary ways in which nitrogen is lost. Denitrification occurs when soil microorganisms use oxygen from the nitrate ion (NO₃⁻) rather than from air in the soil (oxygen) for respiration. This happens most often when soils are saturated and temperatures are warm enough to encourage a high level of microbial activity. Both fertilizer and organic sources of nitrogen are affected. A soil does not have to be completely saturated for denitrification to occur. When it does occur, nitrous oxide gas (N₂O) and other N gases are produced and lost into the atmosphere [3] (Figure 2).




Figure 2

When soils are warm and wet, Nitrate is easily lost as gas. Long period of wet condition can result tremendous fertilizer losses.


Flooded rice cultivation contributing Climate Change through CH₄ and N₂O emission. When soils are warm and wet, Nitrate is easily lost as gas. Long period of wet condition can result tremendous fertilizer losses which economically and environmentally unfriendly. Beside this CH₄ production increases in wet rice cultivation due to anaerobic respiration of the soil

microbial population due to water logged condition and they can't not take oxygen. Therefore organic matter degradation and other plants residues decomposition take place in oxygen limited condition and proportionally produce methane CH₄ (Figure 3).



More greenhouse gas per grain of rice production

Rice production in many regions of the world has accelerated greenhouse gas emission. Rice produces methane gas by methanogenesis process by standing of water in their growing periods. Mostly other gases are also contribute by rice in the form of nitrification and methanogenesis [5]. More carbon dioxide in the atmosphere and rising temperatures cause rice agriculture to release more of the potent greenhouse gas methane (CH₄) for each kilogram of rice it produces, new research published in this week's online edition of Nature Climate Change reveals [6]. Mosier et al reported that "Our results show that rice agriculture becomes less climate friendly as our atmosphere continues to change (Figure 4). Direct seeded rice (DSR) technology is one of the approach mitigating greenhouse gas (GHG) emission.



This is important, because rice paddies are one of the largest human sources of methane, and rice is the world's second-most produced staple crop," said (Dr Kees Jan van Groenigen), Research Fellow at the Botany Department at the School of Natural Sciences, Trinity College Dublin, and lead author of the study [7]. Van Groenigen, along with colleagues from Northern Arizona University and the University of California in Davis, gathered all published research to date from 63 different experiments on

rice paddies, mostly from Asia and North America. The common theme in the experiments was that they measured how rising temperatures and extra carbon dioxide in the atmosphere affect rice yields and the amount of methane that is released by rice paddies [8]. The research team used a technique called meta-analysis, a statistical tool for finding general patterns in a large body of experimental data. "Two strong patterns emerged when analysed all the data: first, more CO₂ boosted emissions of methane from rice paddies, and second, higher temperatures caused a decline in rice yields," explained Professor Bruce Hungate of Northern Arizona University and co-author of the study. Rising temperatures were found to have only small effects on CH₄ emissions, but because they decrease rice yield, they also increase the amount of CH₄ emitted per kilogram of rice. "Together, higher CO₂ concentrations and warmer temperatures predicted for the end of this century will about double the amount of CH₄ emitted per kilogram of rice produced." explained Professor Chris van Kessel of the University of California in Davis and co-author of the study [9]. They reported that global demand for rice will increase further with a growing world population, our results suggest that without additional measures, the total CH₄ emissions from rice agriculture will strongly increase [3]. However, the authors point out that there are several options available to reduce CH₄ emissions from rice agriculture. For instance, management practices such as mid-season drainage and using alternative fertilizers have been shown to reduce CH₄ emissions from rice paddies. Moreover, by switching to more heat tolerant rice cultivars and by adjusting sowing dates, yield declines due to temperature increases can largely be prevented, thereby reducing the effect of warming on CH₄ emissions per yield [10,11]. These findings, together with our own results really stress the need for mitigation and adaptation measures to secure global food supply while at the same time keeping greenhouse gas emissions in check. van Groenigen concluded.

Increased atmospheric CO₂ and rising temperatures are expected to affect rice yields and greenhouse-gas (GHG) emissions from rice paddies [12]. This is important, because rice cultivation is one of the largest human-induced sources of the potent GHG methane (CH₄) and rice is the world's second-most produced staple crop. The need for meeting a growing global food demand argues for assessing GHG emissions from croplands on the basis of yield rather than land area [3]. Such that efforts to reduce GHG emissions take into consideration the consequences for food production [13]. However, it is unclear whether or how the GHG intensity (that is, yield-scaled GHG emissions) of cropping systems will be affected by future atmospheric conditions [3,14,15]. Here we show, using meta-analysis, that increased atmospheric CO₂ (ranging from 550 to 743ppm) and warming (ranging from +0.8 °C to +6 °C) both increase the GHG intensity of rice cultivation. Increased atmospheric CO₂ increased GHG intensity by 31.4%, because CH₄ emissions are stimulated more than rice yields. Warming increased GHG intensity by 11.8% per 1 °C, largely owing to a decrease in yield [16]. This analysis

suggests that rising CO₂ and warming will approximately double the GHG intensity of rice production by the end of the twenty-first century, stressing the need for management practices that optimize rice production while reducing its GHG intensity as the climate continues to change [17-22].

Conclusion

The adverse impact of climate change is not limited to food security and water availability but also uneven rainfall distribution through which intensify floods leads to dampen soil. Various agronomic crop yields declining and most of the minor crops going to vanish and thus the food security problem a rising. Therefore, a well-coordinated program is necessary to create awareness among the society including the policy makers, general citizen & public, various organizations, industrialists and farmers to sustain the food security and the environment, whereas climate change has to be experienced round the world.

References

1. Kim JK, Krishnan HB (2002) Making rice a perfect food: tuning dreams into reality. *J Crop Prod* 5(1-2): 93-130.
2. Imran, Fayaz Ahmad AAK (2015) Phenology of various rice genotypes as affected by different transplanting dates under cold climatic region of khyber pakhtunkhwa-pakistan. *J Environ and E Science* 5(3).
3. Imran (2017) Climate change is a real fact confronting to agricultural productivity. *Int J Environ Sci Nat Res* 3(3): 555613.
4. Osenberg CW, Sarnelle O, Cooper SD, Holt RD (1999) Resolving ecological questions through meta-analysis: Goals, metrics and models. *Ecology* 80(4): 1105-1117.
5. Van Groenigen KJ, Osenberg CW, Hungate BA (2011) Increased soil emissions of potent greenhouse gases under increased atmospheric CO₂. *Nature* 475(7355): 214-216.
6. Cassman KG, Dobermann A, Walters DT, Yang H (2003) Meeting cereal demand while protecting natural resources and improving environmental quality. *Annu Rev Environ Resour* 28: 315-358.
7. Mosier AR, Halvorson AD, Reule CA, Liu XJJ (2006) Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *J Environ Qual* 35(4): 1584-1598.
8. Forster P (2007) In *Climate Change 2007: The Physical Science Basis*. In: Solomon S et al. (Eds.), Cambridge Univ Press, USA, pp. 129-234.
9. Linquist B, van Groenigen KJ, Adviento-Borbe MA, Pittelkow C, van Kessel C (2012) An agronomic assessment of greenhouse gas emissions from major cereal crops. *Glob Change Biol* 18(1): 194-209.
10. Grassini P, Cassman KG (2012) High-yield maize with large net energy yield and small global warming intensity. *Proc Natl Acad USA* 109(4): 1074-1079.
11. Imran, Khan AA, Akhtar K, Zaheer S, Faisal S, et al. (2015) Rice Seedling Characteristics of Various Genotypes Influenced by Different Sowing Dates in Swat-Pakistan. *J Environ and E Science* 5(1): 122-125.
12. Meehl GA (2007) In *Climate Change 2007: The Physical Science Basis*. In: Solomon S, et al. (Eds.), Cambridge Univ Press, USA, pp. 747-845.
13. Van Groenigen JW, Velthof GL, Oenema O, van Groenigen KJ, van Kessel C (2010) Towards an agronomic assessment of N₂O emissions: A case study for arable crops. *Eur J Soil Sci* 61(6): 903-913.
14. Smith P (2007) In *Climate Change 2007: Mitigation*. In: Metz B, et al. (Eds.), Cambridge Univ Press, USA, pp. 4970-540.
15. Maclean JL, Dawe DC, Hardy B, Hettel GP (2002) *Rice Almanac: Source Book for the Most Important Economic Activity on Earth*. (3rd edn), CABI Publishing, England.
16. Lobell DB, Field CB (2007) Global scale climate-crop yield relationships and the impacts of recent warming. *Environ Res Lett* 2: 014002.
17. Ainsworth EA (2008) Rice production in a changing climate: A meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Glob Change Biol* 14(7): 1642-1650.
18. Peng S (2004) Rice yields decline with higher night temperature from global warming. *Proc Natl Acad Sci* 101(27): 9971-9975.
19. Easterling WE, Parry ML, Canziani OF, Palutikof JP, van der Linden PJ, et al. (2007) *Impacts, Adaptation and Vulnerability*. Cambridge Univ Press. USA, pp. 273-313.
20. Imran AA, Inamullah K, Luqman (2015) Weeding stages and their effect on yield and yield components of rice in upper Swat, Pakistan. *Pak J Weed Sci Res* 21(4): 555-563.
21. Imran, Asad Ali K (2015) Effect of transplanting dates on yield and yield components of various rice genotypes in hilly area cold climatic region of khyberpakhtunhwa-Pakistan. *J Bio agric and Care* 5(7).
22. Khan AA, Imran, Shah Inamullah FA, Khan LN (2015) Phenological Traits of Rice as Influenced by Seedling Age and Number of Seedling per Hill under Temperate Region. *J Bio Agric H Care*.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/ARTOAJ.2017.09.55565](https://doi.org/10.19080/ARTOAJ.2017.09.55565)

Your next submission with Juniper Publishers will reach you the below assets

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission

<https://juniperpublishers.com/online-submission.php>