



Effect of Pregnancy on Oxidative Status of Sheep



Musa Suleman Ibrahim^{1*}, Briskila Oliver Ombugadu² and Yusuf Ibrahim Hussaini³

Department of Animal Science, Nasarawa State University, Keffi, Shabu- Lafia Campus, Nigeria

Submission: January 23, 2023; Published: February 21, 2023

*Corresponding author: Musa, Suleman Ibrahim, Department of Animal Science, Nasarawa State University, Keffi, Shabu- Lafia Campus, Nigeria

Abstract

The objective of the study was to determine the effects of pregnancy on oxidative status of sheep. Specific goal of the study was to determine the impact of pregnancy on serum concentration of antioxidant enzymes and the level of lipid peroxidation. Two groups of sheep (10) consisting of pregnant and non-pregnant ewes were used for the study. The pregnant ewes were at third trimester of pregnancy which was confirmed by abdominal palpation and serum progesterone level above 2.2ng/ml. Serum levels of catalase, superoxide dismutase, glutathione peroxidase enzymes and malondialdehyde were determined. The results indicated similar [$p>0.05$] mean values of serum concentration of catalase, superoxide dismutase and glutathione peroxidase. The serum level of malondialdehyde was significantly higher in pregnant ewes compared to non-pregnant animals. It can be concluded that the third trimester of pregnancy is associated with higher serum levels of malondialdehyde.

Introduction

Reactive oxygen species play a critical role in many physiological processes but are detrimental to body cells at high concentration Hohn [1]. Pregnancy is associated with proinflammatory state due to increased activity of monocytes and macrophages resulting in high concentration of oxidative stress biomarkers Ness [2]. Adverse effects of oxidative stress on reproductive performance of sows have been documented, decrease litter size and birth weight were associated with oxidative stress Zhao [3]. Oxidation of proteins can modify their structure thereby causing adverse effects on functions of the protein. Oxidized proteins can also generate antigens that stimulate immune response Halliwell & Whiteman [4]. Increase oxidative stress has been associated with prolonged concentrate feeding in dairy cows, this is attributed to increase rumen endotoxins associated with high concentrate and low neutral detergent fiber content which may stimulate Para inflammatory reactions and reactive oxygen species Golder [5]. Lipid mobilization associated with low energy balance has also been associated with the increase of reactive oxygen species Pedemera [6].

Reactive oxygen species play a critical role in reproductive processes such as ovulation, fertilization, and early embryo development. However, oxidative stress is associated with embryo losses, and it is involved in the pathogenesis of follicular cyst and repeat breeder syndrome in dairy cows Talkdor [7]. Insufficient antioxidant activity can lead to poor pregnancy outcome including fetal growth restrictions. The metabolic changes associated with pregnancy may result in impaired oxidative status leading to adverse effects on pregnancy establishment and maintenance.

The objective of this study, therefore, was to determine the effects of pregnancy on serum concentration of superoxide dismutase, catalase, and glutathione peroxidase enzymes. The impact of pregnancy on lipid peroxidation was also evaluated as indicated by serum level of malondealdehyde.

Materials and Methods

The objective of the study was to determine the effect of pregnancy on oxidative status of sheep. Two groups of sheep [$n=10$] consisting of pregnant and non-pregnant ewes were used for the study. The pregnant ewes were at the third trimester of pregnancy which was confirmed by abdominal palpation and serum progesterone level above 1.2 ng/ml. Blood samples were collected through the jugular vein for determination antioxidant enzymes concentration.

Determination of Serum Antioxidant Enzymes

Serum level of superoxide dismutase was determined by calorimetric method based on the ability of the enzyme to inhibit phenazine methosulphate mediated reduction of nitro blue as described by Weydert [8]. Catalase activity was measured using a combination of optimized enzymatic conditions and spectrophotometric assay of hydrogen peroxide as described by Laszo [9]. Glutathione peroxidase activity was measured using hydrogen peroxide as a substrate as described by Liu [10]. Serum malondialdehyde was determined in terms of thiobarbituric acid reactive substances using spectrophotometer as reported by Surapon [11].

Statistical Analysis

Data obtained was subjected to student t test using statistical tool for agricultural research [STAR]. Means separation was done using least significant difference where significant differences exist.

Results and Discussion

The effects of pregnancy on oxidative status of sheep are shown on Table 1.

Table 1: Effects of pregnancy on serum level of antioxidant enzymes (mean).

Parameters	Pregnant sheep	Non pregnant sheep	Los
Superoxide dismutase (iu/l)	16.40a	15.03a	NS
Catalase (iu/l)	40.36a	40.43a	NS
Glutathione peroxidase (iu/l)	29.43a	29.83a	NS
Malondialdehyde (mmol/l)	1.76b	1.06a	*

Means within same row bearing different superscript are significantly different ($P < 0.05$), Ns = Not significant, * = Significant ($P < 0.05$), Los = Level of Significance

The results indicated similar mean values of catalase, superoxide dismutase and glutathione peroxidase enzymes in both groups. The serum level of malondialdehyde is however, higher [$p > 0.05$] in pregnant ewes compared to non-pregnant ewes. The concentration of malondialdehyde is an established indicator of cellular injury as well as a biomarker of oxidative stress in cells and tissues Simek [12]. Pregnancy is associated with the increase of generation of reactive oxygen species which is linked to oocyte maturation, luteolysis and embryo implantation. Oxidative stress adversely affects placental functions which may lead to poor pregnancy outcome Hussain [13]. This may be attributed to changing metabolic demands associated with pregnancy. Oxidative stress results from disruption of balance between antioxidant system and reactive oxygen species. Decrease plasma antioxidant have been observed in early phase of pregnancy, while total antioxidant capacity improves in the second and third stage of pregnancy Toescu [14]. The findings of this study showed that pregnancy is associated with potential impairment of oxidative status. Increased serum malondialdehyde indicating higher level of lipid peroxidation was recorded in this study. Supplementation with antioxidants is recommended in pregnant ewes.

Conclusion

It can be concluded that serum levels of catalase, superoxide dismutase and glutathione peroxidase enzymes were not influence by pregnancy at third trimester, whereas the serum level of malondialdehyde was elevated indicating higher lipid peroxidation at third trimester of pregnancy in ewes.

References

- Hohn A, Konia J, Grune T (2013) Protein oxidation in aging and the removal of oxidized proteins. *Journal of Proteomics* 92: 132-159.
- Ness RB (2004) The consequences for human reproduction of a robust inflammatory response. *Quarterly Review biology* 79(4): 383-393.
- Zhao Y, Flower WL, Saraiva A, Kim SW (2013) Effect of social ranks and gestation Housing systems on oxidative stress status, reproductive performance, immune status of sows. *Journal of Animal Science* 91(12):5848-5858.
- Halliwell B, Whiteman M (2004) Measuring reactive species and oxidative damage In vivo and in cell culture. *British Journal of Pharmacology* 142(2): 231-255.
- Golder HM, Lean H, Rabiee AR, King R, Celi R (2013) Effects of grain, fructose and histidine feeding on endotoxins and oxidative stress markers in dairy heifers. *Journal of Dairy Science* 96(12): 7881-7891.
- Pedenera M, Celi P, Garcia SC, Savin HE, Fulkerson WJ (2010) Effect of diet Energy balance and milk production on oxidative stress in early lactating dairy cows. *Veterinary Journal* 186(3): 352-257.
- Talukder S, Ingenhoff L, Kerrisk KL (2014) Plasma oxidative stress biomarkers and Progesterone profile in dairy cows diagnosed with an ovarian follicular cyst. *Vet Q* 34(2): 113-117.
- Werdert C (2003) Suppression of the malignant phenotype in human pancreatic cancer cells by the over expression of manganese superoxide dismutase. *Molecular Cancer Therapy* 2: 361-369.
- Laszlo G (1991) A simple method for determination of serum catalase activity and revision of reference range. *International Journal of Clinical Chemistry* 196(2-3): 143-151.
- Liu J (2004) Redox regulation of pancreatic cancer cells growth: role of glutathione peroxidase in the suppression of the malignant phenotype. *Human Gene Therapy* 15: 239-250.
- Surapon P (2009) Serum level of malondialdehyde in type diabetes mellitus Thai subjects. *Siriraj Medical Journal* 61(1): 20-23.
- Simsek S, Abdurrauf Y, Armagan U (2006) Determination of serum malondialdehyde levels in sheep naturally infected with *Dicrocoelium dendriticum*. *Saylik Bil Dergisi* 20(3): 217-220.
- Hussain T, Martaza G, Metwally E (2021) Role of oxidative stress and antioxidant Balance in pregnancy. *Mediators of inflammation* p. 11.
- Toescu V, Nuhall SL, Martin D, Dunne F (2002) Oxidative stress and normal Pregnancy. *Clinical Endocrinology* 57(5): 609-613.



This work is licensed under Creative Commons Attribution 4.0 License
DOI: [10.19080/JDVS.2023.15.555914](https://doi.org/10.19080/JDVS.2023.15.555914)

**Your next submission with Juniper Publishers
will reach you the below assets**

- Quality Editorial service
- Swift Peer Review
- Reprints availability
- E-prints Service
- Manuscript Podcast for convenient understanding
- Global attainment for your research
- Manuscript accessibility in different formats
(Pdf, E-pub, Full Text, Audio)
- Unceasing customer service

Track the below URL for one-step submission
<https://juniperpublishers.com/online-submission.php>